

Generic Approximation of functions by their Padé approximants, II

G. Fournodavlos

Abstract

In [5] we proved that generically functions defined in any open set can be approximated by a sequence of their Padé approximants, in the sense of uniform convergence on compacta. In this paper we examine a more particular space, $A^\infty(\Omega)$, and prove that we can obtain similar approximation results with functions smooth on the boundary.

Subject Classification MSC2010: primary 41A21, 30K05 secondary 30B10, 30E10, 30K99, 41A10, 41A20.

Key words: Padé approximant, Taylor series, Baire's theorem, Runge's theorem, generic property.

1. Introduction

On a disc every holomorphic function f can be approximated by the partial sums of its Taylor expansion. It is also true that generically in a simply connected domain every holomorphic function is the uniform on compacta limit of a subsequence of the partial sums of its Taylor expansion. The partial sums are polynomials and thus by the maximum principle we are led to uniform approximation on compact sets with connected complement. If we replace the partial sums by the Padé approximants $[p/q]_f$, which are rational functions with poles, then we obtain approximation on compact sets with arbitrary connectivity ([4], [5]).

In [2] it was proved that generically every entire function can be approximated uniformly on compacta by a subsequence $[p_n/q_n]_f$ of its Padé approximants, provided $p_n \rightarrow +\infty$ and $p_n - q_n \rightarrow +\infty$. In [5] we weakened the previous assumption to $p_n \rightarrow +\infty$ only and we extended the result to any simply connected domain. We also

obtained the same approximation on any open subset of $\mathbb{C}$ (of arbitrary connectivity) under the assumption $p_n \rightarrow +\infty$ and $q_n \rightarrow +\infty$.

In the present paper we obtain similar results using smooth functions; that is, holomorphic functions on Ω such that every derivative $f^{(l)}$ extends continuously on $\overline{\Omega}$ ($f \in A^\infty(\Omega)$). In the case of a domain $\Omega \subseteq \mathbb{C}$ such that $(\mathbb{C} \cup \{\infty\} \setminus \overline{\Omega})$ is connected we obtain the result provided $p_n \rightarrow +\infty$; this result is generic in a subset of $A^\infty(\Omega)$, which is the closure of the set of polynomials, under the natural topology. We do not know in general if this subspace is the whole $A^\infty(\Omega)$. If there exists a finite constant $M < \infty$ so that all points $A, B \in \overline{\Omega}$ can be joined in $\overline{\Omega}$ by a curve Γ with length $|\Gamma| \leq M$, then the polynomials are dense in $A^\infty(\Omega)$ ([11]). In the case $q = 0$ the generic result in the closure of polynomials in $A^\infty(\Omega)$ is known ([9]).

Finally in the general case of an open subset $\Omega \subseteq \mathbb{C}$ we obtain a similar generic result in the closure in $A^\infty(\Omega)$ of holomorphic functions in some varying neighborhood of Ω , provided $p_n \rightarrow +\infty$ and $q_n \rightarrow +\infty$. Our method of proof is based on Baire's Category theorem ([7], [8]) and extends the methods of [4] and [13].

2. Preliminaries

Let $\Omega \subseteq \mathbb{C}$ be an open set and let us consider the set $A^\infty(\Omega) = \{f \in H(\Omega): f^{(l)} \text{ extends continuously on } \overline{\Omega}, l = 0, 1, \dots\}$, where $f^{(l)}, l = 0, 1, \dots$ denote the derivatives of the holomorphic function f .

We define the following metric ρ on $A^\infty(\Omega)$:

$$\rho(f, g) = \sum_{l=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{2^{l+n}} \min\{\|f - g\|_{l,n}, 1\},$$

where $\|f - g\|_{l,n} = \sup_{z \in \overline{\Omega} \cap \Delta(0,n)} |f^{(l)} - g^{(l)}|$, $l = 0, 1, \dots$. It is easy to see that a sequence in

$A^\infty(\Omega)$, $(f_m)_{m \in \mathbb{N}}$ converges $f_m \xrightarrow{\rho} f \in A^\infty(\Omega)$, if and only if $f_m^{(l)} \rightarrow f^{(l)}$ uniformly on each compact subset of $\overline{\Omega}$, for every l . The space $(A^\infty(\Omega), \rho)$ is complete.

Let f be a function holomorphic in a neighborhood of 0 and let $f(z) = \sum_{v=0}^{\infty} a_v z^v$ its Taylor series. A Padé approximant $[p/q]_f$ of f , $p, q \in \{0, 1, 2, \dots\}$, is a rational function of the form

$$\frac{\sum_{v=0}^p n_v z^v}{\sum_{v=0}^q d_v z^v}, \quad d_0 = 1.$$

such that its Taylor series $\sum_{v=0}^{\infty} b_v z^v$ coincides with $\sum_{v=0}^{\infty} a_v z^v$ up to the first $p+q+1$ terms; that is $b_v = a_v$ for $v = 0, \dots, p+q$ ([1]).

We notice that in case of $q = 0$ there exists always a unique Padé approximant of f and $[p/q]_f(z) = S_p(z)$, where $S_p(z) = \sum_{v=0}^p a_v z^v$. For $q \geq 1$ it is true that there exists a unique Padé approximant of f , if and only if the following determinant is not zero:

$$\det \begin{vmatrix} a_{p-q+1} & a_{p-q+2} & \cdots & a_p \\ a_{p-q+2} & a_{p-q+3} & \cdots & a_{p+1} \\ \vdots & \vdots & & \vdots \\ a_p & a_{p+1} & \cdots & a_{p+q-1} \end{vmatrix} \neq 0, \quad a_i = 0, \quad \text{when } i < 0. \quad (*)$$

Then we write $f \in D_{p,q}$.

If $f \in D_{p,q}$, then $[p/q]_f$ ($q \geq 1$) is given by the Jacobi explicit formula:

$$[p/q]_f = \frac{\det \begin{vmatrix} z^q S_{p-q}(z) & z^{q-1} S_{p-q+1} & \cdots & S_p(z) \\ a_{p-q+1} & a_{p-q+2} & \cdots & a_{p+1} \\ \vdots & \vdots & & \vdots \\ a_p & a_{p+1} & \cdots & a_{p+q} \end{vmatrix}}{\det \begin{vmatrix} z^q & z^{q-1} & \cdots & 1 \\ a_{p-q+1} & a_{p-q+2} & \cdots & a_{p+1} \\ \vdots & \vdots & & \vdots \\ a_p & a_{p+1} & \cdots & a_{p+q} \end{vmatrix}},$$

$$\text{with } S_k(z) = \begin{cases} \sum_{v=0}^k a_v z^v, & k \geq 0 \\ 0, & k < 0. \end{cases}$$

Remark 2.1. If all of the coefficients $\frac{f^{(v)}(0)}{v!} = a_v$, $v = 0, 1, \dots, p+q$, involved in the determinant (*) depend linearly on $d \in \mathbb{C}$, $a_v = c_v \cdot d + \tau_v$, such that $c_v = 0$, when $v < p$ and $c_p \neq 0$, then the determinant is a polynomial in d of degree q and hence only for finite values of d the determinant is zero.

If L is any set we write $h \in H(L)$ if h is holomorphic in some open set containing L . We also denote $\|h\|_L = \sup_{z \in L} |h(z)|$, for every function $h : L \rightarrow \mathbb{C}$ on the set L .

Lemma 2.2. Let $r > 0$, $p, q, s \in \mathbb{N}$ and $K \subseteq \mathbb{C}$ a compact set. If $f \in H(\overline{\Delta(0, r)})$, $f \in D_{p,q}$ such that its Padé approximant $[p/q]_f$ has no poles in K , then for every $\varepsilon > 0$ there exists $\delta > 0$ such that for every $g \in H(\overline{\Delta(0, r)})$ with $\|g - f\|_{\overline{\Delta(0, r)}} < \delta$ it holds $g \in D_{p,q}$ and $\|[p/q]_g^{(l)} - [p/q]_f^{(l)}\|_K < \varepsilon$, $\forall l \in \{0, 1, \dots, s\}$.

Proof. Let $\varepsilon > 0$. Observe that the determinant $(*)$ and the coefficients of the numerator and the denominator of $[p/q]_f^{(l)}$, $l = 0, 1, \dots, s$ depend polynomially on $\frac{f^{(v)}(0)}{v!}$, $v = 0, 1, \dots, p+q$. This implies that there exists $\tilde{\delta}$ such that for every $g \in H(\overline{\Delta(0, r)})$ with $|\frac{g^{(v)}(0)}{v!} - \frac{f^{(v)}(0)}{v!}| < \tilde{\delta}$, $v = 0, 1, \dots, p+q$ it holds $g \in D_{p,q}$ and $\|[p/q]_g^{(l)} - [p/q]_f^{(l)}\|_K < \varepsilon$, $l = 0, 1, \dots, s$.

If $0 < \delta < \min\{r^v \cdot \tilde{\delta} \mid v = 0, 1, \dots, p+q\}$ and $\|g - f\|_{\overline{\Delta(0, r)}} < \delta$, then by Cauchy's estimates we obtain:

$$\left| \frac{g^{(v)}(0)}{v!} - \frac{f^{(v)}(0)}{v!} \right| = \left| \frac{(g - f)^{(v)}(0)}{v!} \right| \leq \frac{\|g - f\|_{\overline{\Delta(0, r)}}}{r^v} < \frac{\delta}{r^v} < \tilde{\delta}. \quad \blacksquare$$

Remark 2.3. It follows from Lemma 2.2 that $D_{p,q} \cap A^\infty(\Omega)$ is open ($0 \in \Omega$).

3. A special case

Let $\Omega \subseteq \mathbb{C}$ be an open set containing 0, such that $(\mathbb{C} \cup \infty) \setminus \overline{\Omega}$ is connected. Also, let $F \subseteq \mathbb{N} \times \mathbb{N}$ which contains a sequence $(\tilde{p}_m, \tilde{q}_m)_{m \in \mathbb{N}}$, such that $\tilde{p}_m \rightarrow +\infty$. We define

- $B_F = \{f \in A^\infty(\Omega) : \text{there exists } (p_m, q_m)_{m \in \mathbb{N}} \text{ in } F \text{ such that } f \in D_{p_m, q_m}, \text{ for all } m \in \mathbb{N} \text{ and for every } K \subseteq \overline{\Omega} \text{ compact } [p_m/q_m]_f^{(l)} \rightarrow f^{(l)} \text{ uniformly on } K, \text{ for each } l = 0, 1, \dots\}$.
- $E(n, s, (p, q)) = \{f \in A^\infty(\Omega) : f \in D_{p,q} \text{ and } \|[p/q]_f - f\|_{l,n} < 1/s, l = 0, 1, \dots, s\}$, $n, s \in \mathbb{N}, (p, q) \in F$.

Lemma 3.1. $B_F = \bigcap_{n,s=1}^{\infty} \bigcup_{(p,q) \in F} E(n, s, (p, q))$.

Proof. It is standard and is omitted. [A similar proof can be found in [12]]. $\blacksquare$

Lemma 3.2. $E(n, s, (p, q))$ is open.

Proof. $D_{p,q} \cap A^\infty(\Omega)$ is open (Remark 2.3) and similarly to the proof of the Lemma 2.2, we can prove that the map $f \mapsto \|[p/q]_f - f\|_{l,n}$ is continuous, for any l . $\blacksquare$

We will now focus our attention on a more accessible space, $H(\overline{\Omega})$, which is a subspace of $A^\infty(\Omega)$ and is considered with its relative topology.

Lemma 3.3. The polynomials are dense in $H(\overline{\Omega})$.

Proof. Let $f \in H(\overline{\Omega})$ and $\varepsilon > 0$. It suffices to show that for $N = N(\varepsilon) \in \mathbb{N}$ and $L = L(\varepsilon) \in \mathbb{N}$ there exists a polynomial P such that $\|P - f\|_{l,N} < \varepsilon$, $\forall l \leq L$. Observe that $(\mathbb{C} \cup \infty) \setminus (\overline{\Omega} \cap \overline{\Delta(0, N)})$ is connected.

- $f \in H(\overline{\Omega})$, thus, there exists $U \subseteq \mathbb{C}$ open such that $f \in H(U)$ and $\overline{\Omega} \subseteq U$.
- It is true that we can find $V \subseteq \mathbb{C}$ open, such that $\overline{\Omega} \cap \overline{\Delta(0, N)} \subseteq V \subseteq U$ and $(\mathbb{C} \cup \infty) \setminus V$ connected (in other words V is simply connected) ([3], [6]).

By Runge's theorem there exists a sequence of polynomials $(P_i)_{i \in \mathbb{N}}$, such that $P_i^{(l)} \rightarrow f^{(l)}$ uniformly on each compact subset of V for every l , which completes the proof. ■

Theorem 3.4. $B_F \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is G_δ and dense in $cl_{A^\infty(\Omega)} H(\overline{\Omega})$. (Hence $B_F \neq \emptyset$).

Proof. Lemma 3.2 implies that $\bigcup_{(p,q) \in F} E(n, s(p, q)) \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is open in $cl_{A^\infty(\Omega)} H(\overline{\Omega})$.

By Lemma 3.1 $B_F \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is G_δ in $cl_{A^\infty(\Omega)} H(\overline{\Omega})$. We claim that $\bigcup_{(p,q) \in F} E(n, s, (p, q)) \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is dense in $cl_{A^\infty(\Omega)} H(\overline{\Omega})$. If this is true, then Baire's Category theorem completes the proof. By Lemma 3.3 it suffices to prove that for every polynomial P and $\varepsilon > 0$ there exists $f \in \bigcup_{(p,q) \in F} E(n, s, (p, q)) \cap H(\overline{\Omega})$ such that $\|P - f\|_{l,N} < \varepsilon$, for every $l \leq L = L(\varepsilon) \in \mathbb{N}$, where $N = N(\varepsilon) \in \mathbb{N}$.

- Let P be a polynomial and $\varepsilon > 0$. There exists $(p, q) \in F$ such that $p > \deg P$.

If $q = 0$, define $f(z) = P(z) + dz^p$, $d \in \mathbb{C} \setminus \{0\}$. It is immediate that $f \in D_{p,q}$ and $[p/q]_f = f$. It follows $f \in E(n, s, (p, q)) \cap H(\overline{\Omega})$. In addition, $\|f - P\|_{l,N} = |d| \cdot \|z^p\|_{l,N} < \varepsilon$, $\forall l \leq L$, when $0 < |d| < \varepsilon / \max_{0 \leq l \leq L} \|z^p\|_{l,N}$.

If $q \geq 1$, we define $\tilde{f}_j(z) = \frac{P(z) + d_j z^p}{1 - (c_j z)^q}$, $c_j, d_j \in \mathbb{C} \setminus \{0\}$, where d_j and c_j will be determined later on, $j \in \mathbb{N}$.

- Let $\lambda > \max\{n, N\}$. We have $\inf_{z \in \Delta(0, \lambda)} |1 - (c_j z)^q| \geq 1 - |c_j|^q \cdot \|z\|_{\Delta(0, \lambda)}^q > \frac{1}{2}$, when $0 < |c_j| < \frac{1}{2^{\frac{1}{q}} \|z\|_{\Delta(0, \lambda)}^q}$, $j \in \mathbb{N}$.
- We have $\|\tilde{f}_j(z) - P(z)\|_{\Delta(0, \lambda)} = \left\| \frac{P(z) + d_j z^p}{1 - (c_j z)^q} \right\|_{\Delta(0, \lambda)} \leq 2(\|P(z)\|_{\Delta(0, \lambda)} + |c_j|^q \cdot \|z\|_{\Delta(0, \lambda)}^q + |d_j| \cdot \|z\|_{\Delta(0, \lambda)}^q)$. Thus, there exists $\delta_j > 0$, $j \in \mathbb{N}$, such that: $\|\tilde{f}_j - P\|_{\Delta(0, \lambda)} < 1/j$, when $0 < |c_j| < \delta_j < \frac{1}{2^{\frac{1}{q}} \|z\|_{\Delta(0, \lambda)}^q}$ and $0 < |d_j| < \delta_j$. Hence, $\tilde{f}_j \rightarrow P$ uniformly on $\overline{\Delta(0, \lambda)}$ and so $\tilde{f}_j^{(l)} \rightarrow P^{(l)}$ uniformly on $\overline{\Omega} \cap \overline{\Delta(0, N)}$ (which is contained in $\Delta(0, \lambda)$), for every l . Therefore, there exists $j_0 \in \mathbb{N}$ such that $\|\tilde{f}_{j_0} - P\|_{l,N} < \varepsilon/2$, $l = 0, 1, \dots, L$.
- We fix c_{j_0} satisfying the above. Around 0, $\tilde{f}_{j_0}(z) = P(z) + d_{j_0} z^p + P(z) \cdot (c_{j_0} z)^q + d_{j_0} z^p \cdot (c_{j_0} z)^q + \dots$. According to Remark 2.1 we can choose $0 < |d_{j_0}| < \delta_{j_0}$, such

that $\tilde{f}_{j_0} \in D_{p,q}$. By the uniqueness of the Padé approximant of $\tilde{f}_{j_0}$ we obtain $[p/q]_{\tilde{f}_{j_0}} = \tilde{f}_{j_0}$.

- Let $r > 0$: $\overline{\Delta(0, r)} \subseteq \Omega \cap \Delta(0, \lambda)$. By Lemma 2.2 there exists $\delta > 0$ such that for every $f \in H(\overline{\Delta(0, r)})$ with $\|f - \tilde{f}_{j_0}\|_{\overline{\Delta(0, r)}} < \delta$ it holds $\|[p/q]_f - [p/q]_{\tilde{f}_{j_0}}\|_{l,n} < 1/2s$, $\forall l \leq s$. Also, we demand $0 < \delta < \min\{1/2s, \varepsilon/2\}$.
- The Taylor series of $\tilde{f}_{j_0}$ around 0 has radius of convergence greater than $\frac{1}{|c_{j_0}|} > \lambda$. It follows that its partial sums $(\sum_{v=0}^k \frac{\tilde{f}_{j_0}^{(v)}(0)}{v!} z^v)^{(l)} \rightarrow \tilde{f}_{j_0}^{(l)}$ uniformly on $\overline{\Omega} \cap \overline{\Delta(0, \lambda)}$, for every l . Hence, there exists a partial sum $f(z) = \sum_{v=0}^{k_0} \frac{\tilde{f}_{j_0}^{(v)}(0)}{v!} z^v$ such that $\|f - \tilde{f}_{j_0}\|_{l,\lambda} < \delta$, $\forall l \leq \max\{s, L\}$.
- f satisfies: $\|[p/q]_f - f\|_{l,n} \leq \|[p/q]_f - [p/q]_{\tilde{f}_{j_0}}\|_{l,n} + \|\tilde{f}_{j_0} - f\|_{l,n} < 1/2s + \delta < 1/s$, $\forall l \leq s$. It follows that $f \in E(n, s, (p, q)) \cap H(\overline{\Omega})$. Also, it holds $\|f - P\|_{l,N} \leq \|f - \tilde{f}_{j_0}\|_{l,N} + \|\tilde{f}_{j_0} - P\|_{l,N} < \varepsilon/2 + \delta < \varepsilon$, $\forall l \leq L$.

This completes the proof. ■

4. The general case

Let $\Omega \subseteq \mathbb{C}$ be an open set containing 0. Also, let $F \subseteq \mathbb{N} \times \mathbb{N}$ which contains a sequence $(\tilde{p}_m, \tilde{q}_m)_{m \in \mathbb{N}}$ such that $\tilde{p}_m \rightarrow +\infty$ and $\tilde{q}_m \rightarrow +\infty$. We define B_F and $E(n, s, (p, q))$ similarly as in Section 3.

The analogue of Lemmas 3.1, 3.2 hold in this case also. Like before we concentrate on $H(\overline{\Omega})$ and its closure in $A^\infty(\Omega)$.

Lemma 4.1. *The rational functions with poles off $\overline{\Omega}$ are dense in $H(\overline{\Omega})$.*

Proof. Let $f \in H(\overline{\Omega})$. There exists $U \subseteq \mathbb{C}$ open (depending on f) such that $\overline{\Omega} \subseteq U$ and $f \in H(U)$. By Runge's theorem there exists a sequence $(R_i)_{i \in \mathbb{N}}$ of rational functions with poles in $(\mathbb{C} \cup \infty) \setminus U$, hence $R_i \in H(U) \subseteq H(\overline{\Omega})$, $\forall i \in \mathbb{N}$, such that $R_i \rightarrow f$ uniformly on each compact set of U . Similarly to the Lemma 3.3, for a given ε there exists i_0 such that $\|R_i - f\|_{l,N} < \varepsilon/2$, $\forall l \leq L$, where $L = L(\varepsilon) \in \mathbb{N}$, $N = N(\varepsilon) \in \mathbb{N}$ are chosen so that $\rho(f, R_{i_0}) < \varepsilon$. ■

Theorem 4.2. *$B_F \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is G_δ and dense in $cl_{A^\infty(\Omega)} H(\overline{\Omega})$. (Hence $B_F \neq \emptyset$).*

Proof. Since $\bigcup_{(p,q) \in F} E(n, s, (p, q))$ is open, it follows that $B_F \cap cl_{A^\infty(\Omega)} H(\overline{\Omega})$ is G_δ in the subspace. By Baire's Category theorem the proof would be complete if the set

$\bigcup_{(p,q) \in F} E(n, s, (p, q)) \cap H(\overline{\Omega})$ was dense in $H(\overline{\Omega})$, $n, s \in \mathbb{N}$. By Lemma 4.1 it suffices to show that for every rational function R with poles off $\overline{\Omega}$ (or $R \in H(\overline{\Omega})$) and every $\varepsilon > 0$ there exists $f \in \bigcup_{(p,q) \in F} E(n, s, (p, q)) \cap H(\overline{\Omega})$ such that $\|f - R\|_{l,N} < \varepsilon$, $\forall l \leq L = L(\varepsilon) \in \mathbb{N}$, $N = N(\varepsilon) \in \mathbb{N}$.

- Let $R(z) = \frac{A(z)}{B(z)}$ be a rational function with poles only in $(\mathbb{C} \cup \{\infty\}) \setminus \overline{\Omega}$, where A, B are polynomials and let $\varepsilon > 0$. There exists $(p, q) \in F$ such that $p > \deg A$ and $q > \deg B$. We define $\tilde{f}_j(z) = \frac{A(z) + d_j z^p}{B(z) - (c_j z)^q}$, $c_j, d_j \in \mathbb{C} \setminus \{0\}$, where c_j and d_j will be determined later on, $j \in \mathbb{N}$.

- Since R has no poles in $\overline{\Omega}$, there exists $U \subseteq \mathbb{C}$ open such that $R \in H(U)$ and $\overline{\Omega} \subseteq U$. Also, there exists $K \subseteq U$ compact such that $K^0 \supseteq \overline{\Omega} \cap \overline{\Delta(0, \lambda)}$, where $\lambda = \max\{n, N\}$, and every component of $(\mathbb{C} \cup \infty) \setminus K$ contains at least one component of $(\mathbb{C} \cup \infty) \setminus U$ ([14]).

- We have $B(0) \neq 0$ and $\inf_{z \in K} |B(z)| > 0$. Furthermore, $\inf_{z \in K} |B(z) - (c_j z)^q| \geq \inf_{z \in K} |B(z)| - |c_j|^q \cdot \|z\|_K^q > 0$, when $0 < |c_j| < \left(\frac{\inf_K |B(z)|}{\|z\|_K^q}\right)^{1/q}$, $\forall j \in \mathbb{N}$. Thus,

$$\|\tilde{f}_j(z) - R(z)\|_K = \left\| \frac{A(z)(c_j z)^q + B(z)d_j z^p}{B(z)(B(z) - (c_j z)^q)} \right\|_K \leq \frac{\|A(z)\|_K \cdot |c_j|^q \cdot \|z\|_K^q + \|B(z)\|_K \cdot |d_j| \cdot \|z\|_K^p}{\inf_{z \in K} |B(z)| \cdot \inf_{z \in K} |B(z) - (c_j z)^q|}.$$

- There exists $\delta_j > 0$, $j \in \mathbb{N}$, such that $\|\tilde{f}_j - R\|_K < 1/j$, whenever $|c_j| < \delta_j < \left(\frac{\inf_K |B(z)|}{\|z\|_K^q}\right)^{1/q}$ and $|d_j| < \delta_j$, $\forall j \in \mathbb{N}$. Hence, $\tilde{f}_j \rightarrow R$ uniformly on K and $\tilde{f}_j^{(l)} \rightarrow R^{(l)}$ on each compact subset of K^0 , for every l . This implies that there exists $j_0 \in \mathbb{N}$, such that $\|\tilde{f}_{j_0} - R\|_{l,N} < \varepsilon/2$, $\forall l \leq L$.

- We fix c_{j_0} satisfying the above. Around 0 we have: $\tilde{f}_{j_0}(z) = B^{-1}(0)A(z) + B^{-1}(0)d_{j_0}z^p - B^{-1}(0)A(z) \cdot (\tilde{B}(z) - 1) - B^{-1}(0)d_{j_0}z^p \cdot (\tilde{B}(z) - 1) + \dots$, where $\tilde{B}(z) = B^{-1}(0)B(z) - B^{-1}(0)(c_{j_0}z)^q$. By Remark 2.1 we can choose $0 < |d_{j_0}| < \delta_{j_0}$ such that $\tilde{f}_{j_0} \in D_{p,q}$. Thus, there exists a unique Padé approximant of $\tilde{f}_{j_0}$ and $\tilde{f}_{j_0} = \frac{B^{-1}(0)A(z) + B^{-1}(0)d_{j_0}z^p}{B^{-1}(0)B(z) - B^{-1}(0)(c_{j_0}z)^q}$ satisfies $[p/q]_{\tilde{f}_{j_0}} = \tilde{f}_{j_0}$.

- There exists $r > 0$: $\overline{\Delta(0, r)} \subseteq \Omega \cap \Delta(0, \lambda) \subseteq K^0$. Lemma 2.2 provides $0 < \delta < \min\{1/2s, \varepsilon/2\}$ such that for every $f \in H(\overline{\Delta(0, r)})$ with $\|f - \tilde{f}_{j_0}\|_{\overline{\Delta(0, r)}} < \delta$, it follows $f \in D_{p,q}$ and $\|[p/q]_f - [p/q]_{\tilde{f}_{j_0}}\|_{l,n} < 1/2s$, $\forall l \leq s$.

- By Runge's theorem there exists a sequence of rational functions, $(R_i)_{i \in \mathbb{N}}$, with poles off K and more particularly (see previous property of K) off U , such that

$R_i \rightarrow \tilde{f}_{j_0}$ uniformly on K . This implies that $R_i^{(l)} \rightarrow \tilde{f}_{j_0}^{(l)}$ uniformly on each compact subset of K^0 . Hence, there exists $f = R_{i_0} \in H(U) \subseteq H(\overline{\Omega})$ such that $\|f - \tilde{f}_{j_0}\|_{l,\lambda} < \delta$, $\forall l \leq \max\{s, L\}$, because $\overline{\Omega} \cap \overline{\Delta(0, \lambda)} \subseteq K^0$.

- It follows that $\|[p/q]_f - f\|_{l,n} \leq \|[p/q]_f - [p/q]_{\tilde{f}_{j_0}}\|_{l,n} + \|\tilde{f}_{j_0} - f\|_{l,n} < 1/2s + \delta < 1/s$, $\forall l \leq s$. Thus, $f \in E(n, s, (p, q)) \cap H(\overline{\Omega})$. Moreover, it holds $\|f - R\|_{l,N} \leq \|f - \tilde{f}_{j_0}\|_{l,N} + \|\tilde{f}_{j_0} - R\|_{l,N} < \delta + \varepsilon/2 < \varepsilon$, $\forall l \leq L$.

This completes the proof. ■

References

- [1] G. A. Baker, Jr. and P. R. Graves-Morris: Padé Approximants, Vol. 1 and 2, (Encyclopedia of Mathematics and its Applications), Cambridge Un. Press, 2010.
- [2] P. B. Borwein: The usual behaviour of rational approximants, Canad. Math. Bull. Vol. 26, 1983, p. 317-323.
- [3] G. Costakis: Some remarks on universal functions and Taylor series, Math. Proc. Cambr. Philos. Soc., Vol. 128, 2000, p. 157-175.
- [4] N. J. Daras and V. Nestoridis: Universal Padé approximation, arXiv: 1102.4782v1, 2011.
- [5] G. Fournodavlos: Generic Approximation of functions by their Padé approximants I, arXiv: 1103.3841, 2011.
- [6] K.-G. Grosse-Erdmann: Holomorphe Monster und universelle Funktionen, Mitt. Math. Sem. Giessen, Vol. 176, 1987, p. 1-84.
- [7] K.-G. Grosse-Erdmann: Universal families and hypercyclic operators, Bull. Amer. math. Soc., Vol. 36, 1999, p. 345-381.
- [8] J.-P. Kahane: Baire's Category theorem and trigonometric series, J. Anal. Math., Vol. 80, 2000, p. 143-182.
- [9] Ch. Kariofillis, Ch. Konstandaki and V. Nestoridis: Smooth universal Taylor series, Monatsh. Math., Vol. 147, Issue 3, 2006, p. 249-257.
- [10] A. Melas and V. Nestoridis: Universality of Taylor series as a generic property of holomorphic functions, Adv. Math. Vol. 157, 2001, p. 138-176.

- [11] A. Melas and V. Nestoridis: On various types of universal Taylor series, Complex Variables Theory, Vol. 44, Issue 3, 2001, p. 245-258.
- [12] V. Nestoridis: Universal Taylor series, Annales de l' Institute Fourier, Vol. 46, 1996, p. 1293-1306.
- [13] V. Nestoridis: A strong notion of universal Taylor series, J. London Math. Soc., Vol. 68, Issue 2, 2003, p. 712-724.
- [14] W. Rudin, Real and Complex Analysis, McGraw-Hill, 1986.

Department of Mathematics,
University of Athens
Panepistemiopolis, 157 84 Athens, Greece
e-mail: gregdavlos@hotmail.com